



ICD-9-CM Code	ICD-10-CM Code(s)	
<b>789.00</b>	Abdominal pain, unspecified site	
ICD-10-CM Codes	<b>R10.9</b> Unspecified abdominal pain <b>*There</b> are more specific code choice selections available in ICD-10-CM. These include: <b>R10.0</b> Acute abdomen <b>R10.10</b> Upper abdominal pain, unspecified <b>R10.11</b> Right upper quadrant pain <b>R10.12</b> Left upper quadrant pain <b>R10.13</b> Epigastric pain <b>R10.2</b> Pelvic and perineal pain <b>R10.30</b> Lower abdominal pain, unspecified <b>R10.31</b> Right lower quadrant pain <b>R10.32</b> Left lower quadrant pain <b>R10.33</b> Periumbilical pain <b>R10.84</b> Generalized abdominal pain	
<b>461.9</b>	Acute sinusitis, unspecified	
ICD-10-CM Codes	<b>J01.90</b> Acute sinusitis, unspecified <b>**Use additional code (B95-B97)</b> to identify infectious agent <b>*There</b> are more specific code choice selections available in ICD-10-CM. These include: <b>J01.00</b> Acute maxillary sinusitis, unspecified <b>J01.10</b> Acute frontal sinusitis, unspecified <b>J01.20</b> Acute ethmoidal sinusitis, unspecified <b>J01.30</b> Acute sphenoidal sinusitis, unspecified <b>J01.40</b> Acute pansinusitis, unspecified <b>J01.80</b> Other acute sinusitis	
<b>784.0</b>	Headache	<b>R51</b> Headache
<b>250.00</b>	Diabetes mellitus without mention of complications type II or unspecified type, not stated as uncontrolled	<b>E11.9</b> Type 2 diabetes mellitus without complications
<b>599.0</b>	Urinary tract infection, site not specified	
ICD-10-CM Codes	<b>N39.0</b> Urinary tract infection, site not specified <b>** Use additional code (B95-B97)</b> to identify infectious agent <b>** There</b> are more specific code choice selections available in ICD-10-CM. These include: <b>N30.00</b> Acute cystitis without hematuria <b>N30.01</b> Acute cystitis with hematuria <b>N30.10</b> Interstitial cystitis (chronic) without hematuria <b>N30.11</b> Interstitial cystitis (chronic) with hematuria <b>N30.20</b> Other chronic cystitis without hematuria <b>N30.21</b> Other chronic cystitis with hematuria <b>N30.30</b> Trigonitis without hematuria <b>N30.31</b> Trigonitis with hematuria <b>N30.40</b> Irradiation cystitis without hematuria <b>N30.41</b> Irradiation cystitis with hematuria <b>N30.80</b> Other cystitis without hematuria <b>N30.81</b> Other cystitis with hematuria	
<b>719.41</b>	Pain in joint, shoulder region	
ICD-10-CM Codes	<b>M25.511</b> Pain in right shoulder <b>M25.512</b> Pain in left shoulder <b>M25.519</b> Pain in unspecified shoulder	

ICD-9-CM Code	ICD-10-CM Code(s)	
<b>311</b>	Depressive disorder, not elsewhere classified	
ICD-10-CM Codes	<b>F32.9</b> Major depressive disorder, single episode, unspecified <b>*There</b> are more specific code choice selections available in ICD-10-CM. These include: <b>F32.0</b> Major depressive disorder, single episode, mild <b>F32.1</b> Major depressive disorder, single episode, moderate <b>F32.2</b> Major depressive disorder, single episode, severe without psychotic features <b>F32.3</b> Major depressive disorder, single episode, severe with psychotic features <b>F32.4</b> Major depressive disorder, single episode, in partial remission <b>F32.5</b> Major depressive disorder, single episode, in full remission <b>F32.8</b> Other depressive episodes	
<b>466.0</b>	Acute bronchitis	
ICD-10-CM Codes	<b>J20.9</b> Acute bronchitis, unspecified <b>* There</b> are more specific code choice selections available in ICD-10-CM. These include: <b>J20.0</b> Acute bronchitis due to Mycoplasma pneumoniae <b>J20.1</b> Acute bronchitis due to Hemophilus influenzae <b>J20.2</b> Acute bronchitis due to streptococcus <b>J20.3</b> Acute bronchitis due to coxsackievirus <b>J20.4</b> Acute bronchitis due to parainfluenza virus <b>J20.5</b> Acute bronchitis due to respiratory syncytial virus <b>J20.6</b> Acute bronchitis due to rhinovirus <b>J20.7</b> Acute bronchitis due to echovirus <b>J20.8</b> Acute bronchitis due to other specified organisms	
<b>724.5</b>	Backache, unspecified	
ICD-10-CM Codes	<b>M54.9</b> Dorsalgia, unspecified <b>*There</b> are more specific code choice selections available in ICD-10-CM. These include: <b>M54.89</b> Other dorsalgia <b>M54.6</b> Pain in thoracic spine <b>M54.5</b> Low back pain <b>M53.3</b> Sacrococcygeal disorders, not elsewhere classified	
<b>V72.31</b>	Routine gynecological exam	
ICD-10-CM Codes	<b>Z01.411</b> Encounter for gynecological examination (general) (routine) with abnormal findings <b>* Use additional code to identify abnormal findings</b> <b>Z01.419</b> Encounter for gynecological examination (general) (routine) without abnormal findings Instructional note under codes should be as follows: <b>**Use additional code:</b> for screening for human papillomavirus, if applicable ( <b>Z11.51</b> ) for screening vaginal pap smear, if applicable ( <b>Z12.72</b> ) to identify acquired absence of uterus, if applicable ( <b>Z90.71</b> -)	
<b>616.10</b>	Vaginitis and vulvovaginitis, unspecified	
ICD-10-CM Codes	<b>N76.0</b> Acute vaginitis <b>**Use additional code (B95-B97),</b> to identify infectious agent <b>There</b> are other codes from this subcategory to describe vaginitis: <b>N76.1</b> Subacute and chronic vaginitis <b>N76.2</b> Acute vulvitis <b>N76.3</b> Subacute and chronic vulvitis <b>N76.89</b> Other specified inflammation of vagina and vulva	





ICD-9-CM Code	ICD-10-CM Code(s)
<b>V10.83</b>	Personal history of other malignant neoplasm of skin
ICD-10-CM Codes	<b>Z85.828</b> Personal history of other malignant neoplasm of skin *There are more specific code choice selections available in ICD-10-CM. These include: <b>Z85.820</b> Personal history of malignant melanoma of skin <b>Z85.821</b> Personal history of Merkel cell carcinoma
<b>706.2</b>	Sebaceous cyst <b>L72.3</b> Sebaceous cyst
<b>707.9</b>	Chronic ulcer of unspecified site
ICD-10-CM Codes	<b>L98.499</b> Non-pressure chronic ulcer of skin of other sites with unspecified severity *There are more specific code choice selections available in ICD-10-CM. These include: <b>L98.411</b> Non-pressure chronic ulcer of buttock limited to breakdown of skin <b>L98.412</b> Non-pressure chronic ulcer of buttock with fat layer exposed <b>L98.413</b> Non-pressure chronic ulcer of buttock with necrosis of muscle <b>L98.414</b> Non-pressure chronic ulcer of buttock with necrosis of bone <b>L98.419</b> Non-pressure chronic ulcer of buttock with unspecified severity <b>L98.421</b> Non-pressure chronic ulcer of back limited to breakdown of skin <b>L98.422</b> Non-pressure chronic ulcer of back with fat layer exposed <b>L98.423</b> Non-pressure chronic ulcer of back with necrosis of muscle <b>L98.424</b> Non-pressure chronic ulcer of back with necrosis of bone <b>L98.429</b> Non-pressure chronic ulcer of back with unspecified severity <b>L98.491</b> Non-pressure chronic ulcer of skin of other sites limited to breakdown of skin <b>L98.492</b> Non-pressure chronic ulcer of skin of other sites with fat layer exposed <b>L98.493</b> Non-pressure chronic ulcer of skin of other sites with necrosis of muscle <b>L98.494</b> Non-pressure chronic ulcer of skin of other sites with necrosis of bone
<b>272.4</b>	Hyperlipidemia, other and unspecified
ICD-10-CM Codes	<b>E78.4</b> Other hyperlipidemia <b>E78.5</b> Hyperlipidemia, unspecified **There are more specific code choice selections available in ICD-10-CM. These include: <b>E78.0</b> Pure hypercholesterolemia <b>E78.1</b> Pure hyperglyceridemia <b>E78.2</b> Mixed hyperlipidemia <b>E78.3</b> Hyperchylomicronemia <b>E78.6</b> Lipoprotein deficiency
<b>786.59</b>	Chest pain, other
ICD-10-CM Codes	<b>R07.89</b> Other chest pain NOS <b>R07.82</b> Intercostal pain

ICD-9-CM Code	ICD-10-CM Code(s)
<b>782.3</b>	Edema
ICD-10-CM Codes	<b>R60.0</b> Localized edema <b>R60.1</b> Generalized edema <b>R60.9</b> Edema, unspecified
<b>786.2</b>	Cough <b>R05</b> Cough
<b>V70.0</b>	Routine general medical examination at a health care facility
ICD-10-CM Codes	<b>Z00.00</b> Encounter for general adult medical examination without abnormal findings <b>Z00.01</b> Encounter for general adult medical examination with abnormal findings ** Use additional code to identify abnormal findings
<b>729.5</b>	Pain in limb
ICD-10-CM Codes	<b>M79.609</b> Pain in unspecified limb ** There are more specific code choice selections available in ICD-10-CM. These include: <b>M79.601</b> Pain in right arm <b>M79.602</b> Pain in left arm <b>M79.603</b> Pain in arm, unspecified <b>M79.604</b> Pain in right leg <b>M79.605</b> Pain in left leg <b>M79.606</b> Pain in leg, unspecified <b>M79.631</b> Pain in right upper arm <b>M79.622</b> Pain in left upper arm <b>M79.629</b> Pain in unspecified upper arm <b>M79.631</b> Pain in right forearm <b>M79.632</b> Pain in left forearm <b>M79.639</b> Pain in unspecified forearm <b>M79.651</b> Pain in right thigh <b>M79.652</b> Pain in left thigh <b>M79.659</b> Pain in unspecified thigh <b>M79.661</b> Pain in right lower leg <b>M79.662</b> Pain in left lower leg <b>M79.669</b> Pain in unspecified lower leg
<b>724.1</b>	Pain in thoracic spine <b>M54.6</b> Pain in thoracic spine
<b>723.0</b>	Spinal stenosis of cervical region <b>M48.02</b> Spinal stenosis, cervical region
<b>724.3</b>	Sciatica
ICD-10-CM Codes	<b>M54.30</b> Sciatica, unspecified site <b>M54.31</b> Sciatica, right side <b>M54.32</b> Sciatica, left side <b>M54.40</b> Lumbago with sciatica, unspecified site <b>M54.41</b> Lumbago with sciatica, right side <b>M54.42</b> Lumbago with sciatica, left side
<b>722.52</b>	Degeneration of lumbar or lumbosacral intervertebral disc
ICD-10-CM Codes	<b>M51.36</b> Other intervertebral disc degeneration, lumbar region <b>M51.37</b> Other intervertebral disc degeneration, lumbosacral region





ICD-9-CM Code	ICD-10-CM Code(s)
<b>592.0</b>	Calculus of kidney
ICD-10-CM Codes	<b>N20.0</b> Calculus of kidney <b>N20.9</b> Urinary calculus; unspecified <b>N20.2</b> Calculus of kidney and ureter
<b>585.9</b>	Chronic kidney disease, unspecified
ICD-10-CM Codes	<b>N18.9</b> Chronic kidney disease, unspecified ** Code first any associated: Diabetic chronic kidney disease ( <b>E08.22</b> , <b>E09.22</b> , <b>E10.22</b> , <b>E11.22</b> , <b>E13.22</b> ) Hypertensive chronic kidney disease ( <b>I12-</b> , <b>I13-</b> ) Use additional code to identify kidney transplant status, if applicable ( <b>Z94.0</b> ) ** There are more specific code choice selections available in ICD-10-CM. These include: <b>N18.1</b> Chronic kidney disease, stage 1 <b>N18.2</b> Chronic kidney disease, stage 2 (mild) <b>N18.3</b> Chronic kidney disease, stage 3 (moderate) <b>N18.4</b> Chronic kidney disease, stage 4 (severe) <b>N18.5</b> Chronic kidney disease, stage 5 <b>N18.6</b> End stage renal disease ** Use additional code to identify dialysis status ( <b>Z99.2</b> )
<b>443.9</b>	Peripheral vascular disease, unspecified
ICD-10-CM Codes	<b>I73.9</b> Peripheral vascular disease, unspecified ** There are more specific code choice selections available. These include: <b>I73.00</b> Raynaud's syndrome without gangrene <b>I73.01</b> Raynaud's syndrome with gangrene <b>I73.1</b> Thromboangiitis obliterans (Buerger's disease) <b>I73.81</b> Erythromelalgia <b>I73.89</b> Other specified peripheral vascular diseases
<b>780.97</b>	Altered mental status
ICD-10-CM Codes	<b>R41.82</b> Altered mental status, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: <b>R41.0</b> Disorientation, unspecified <b>R41.81</b> Age-related cognitive decline <b>R41.840</b> Attention and concentration deficit <b>R41.89</b> Other symptoms and signs involving cognitive functions and awareness
<b>327.23</b>	Obstructive sleep apnea (adult)(pediatric)
ICD-10-CM Codes	<b>G47.33</b> Obstructive sleep apnea (adult)(pediatric) ** Code also any associated underlying condition
<b>385.30</b>	Cholesteatoma, unspecified
ICD-10-CM Codes	<b>H71.90</b> Unspecified cholesteatoma, unspecified ear <b>H71.91</b> Unspecified cholesteatoma, right ear <b>H71.92</b> Unspecified cholesteatoma, left ear <b>H71.93</b> Unspecified cholesteatoma, bilateral ** There are more specific code choice selections available in ICD-10-CM. These include: <b>H71.00</b> Cholesteatoma of attic, unspecified ear <b>H71.01</b> Cholesteatoma of attic, right ear

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes (cont.)	<b>H71.02</b> Cholesteatoma of attic, left ear <b>H71.03</b> Cholesteatoma of attic, bilateral <b>H71.10</b> Cholesteatoma of tympanum, unspecified ear <b>H71.11</b> Cholesteatoma of tympanum, right ear <b>H71.12</b> Cholesteatoma of tympanum, left ear <b>H71.13</b> Cholesteatoma of tympanum, bilateral <b>H71.20</b> Cholesteatoma of mastoid, unspecified ear <b>H71.21</b> Cholesteatoma of mastoid, right ear <b>H71.22</b> Cholesteatoma of mastoid, left ear <b>H71.23</b> Cholesteatoma of mastoid, bilateral <b>H60.40</b> Cholesteatoma of external ear, unspecified ear <b>H60.41</b> Cholesteatoma of right external ear <b>H60.42</b> Cholesteatoma of left external ear <b>H60.43</b> Cholesteatoma of external ear, bilateral
<b>715.00</b>	Osteoarthritis, generalized, site unspecified
ICD-10-CM Codes	<b>M15.0</b> Primary generalized (osteo)arthritis <b>M15.4</b> Erosive (osteo)arthritis <b>M15.8</b> Other polyosteoarthritis <b>M15.9</b> Polyosteoarthritis, unspecified
<b>729.1</b>	Myalgia and myositis, unspecified
ICD-10-CM Codes	<b>M79.1</b> Myalgia <b>M79.7</b> Fibromyalgia <b>M60.9</b> Myositis, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: <b>M60.811</b> Other myositis, right shoulder <b>M60.822</b> Other myositis, left shoulder <b>M60.819</b> Other myositis, unspecified shoulder <b>M60.821</b> Other myositis, right upper arm <b>M60.822</b> Other myositis, left upper arm <b>M60.829</b> Other myositis, unspecified upper arm <b>M60.831</b> Other myositis, right forearm <b>M60.832</b> Other myositis, left forearm <b>M60.839</b> Other myositis, unspecified forearm <b>M60.841</b> Other myositis, right hand <b>M60.842</b> Other myositis, left hand <b>M60.849</b> Other myositis, unspecified hand <b>M60.851</b> Other myositis, right thigh <b>M60.852</b> Other myositis, left thigh <b>M60.859</b> Other myositis, unspecified thigh <b>M60.861</b> Other myositis, right lower leg <b>M60.862</b> Other myositis, left lower leg <b>M60.869</b> Other myositis, unspecified lower leg <b>M60.871</b> Other myositis, right ankle and foot <b>M60.872</b> Other myositis, left ankle and foot <b>M60.879</b> Other myositis, unspecified ankle and foot





ICD-9-CM Code	ICD-10-CM Code(s)
<b>726.5</b>	Enthesopathy of hip region
ICD-10-CM Codes	<p><b>M70.60</b> Trochanteric bursitis, unspecified hip  <b>M70.61</b> Trochanteric bursitis, right hip  <b>M70.62</b> Trochanteric bursitis, left hip  <b>M70.70</b> Other bursitis, unspecified hip  <b>M70.71</b> Other bursitis, right hip  <b>M70.72</b> Other bursitis, left hip                      ** Use additional external cause code to identify activity causing disorder (Y93-)  <b>M76.10</b> Psoas tendinitis, unspecified hip  <b>M76.11</b> Psoas tendinitis, right hip  <b>M76.12</b> Psoas tendinitis, left hip  <b>M76.20</b> Iliac crest spur, unspecified hip  <b>M76.21</b> Iliac crest spur, right hip  <b>M76.22</b> Iliac crest spur, left hip</p>
<b>401.1</b>	Essential hypertension, benign <b>I10</b> Essential (primary) hypertension
<b>600.01</b>	Hypertrophy (benign) of prostate with urinary obstruction and other lower urinary tract symptoms (LUTS) lower urinary tract symptoms (LUTS)
ICD-10-CM Codes	<p>Two codes are required in ICD-10-CM:  <b>N40.1</b> Enlarged prostate with lower urinary tract symptoms (LUTS); AND  <b>N13.8</b> Other obstructive and reflux uropathy                      ** Note under <b>N40.1</b> states – Use additional code for associated symptoms, when specified:                      • Incomplete bladder emptying (<b>R39.14</b>)                      • Nocturia (<b>R35.1</b>)                      • Straining on urination (<b>R39.16</b>)                      • Urinary frequency (<b>R35.0</b>)                      • Urinary hesitancy (<b>R39.11</b>)                      • Urinary incontinence (<b>N39.4-</b>)                      • Urinary obstruction (<b>N13.8</b>)                      • Urinary retention (<b>R33.8</b>)                      • Urinary urgency (<b>R39.15</b>)                      • Weak urinary stream (<b>R39.12</b>)                      ** Note under <b>N13.8</b> states – Code first, if applicable, any causal condition, such as:                      • Enlarged prostate (<b>N40.1</b>)</p>
<b>474.00</b>	Chronic tonsillitis
ICD-10-CM Codes	<p><b>J35.01</b> Chronic tonsillitis                      * Use additional code to identify:                      Exposure to environmental tobacco smoke (<b>Z77.22</b>)                      Exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)                      History of tobacco use (<b>Z87.891</b>)                      Occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)                      Tobacco dependence (<b>F17-</b>)                      Tobacco use (<b>Z72.0</b>)</p>
<b>474.10</b>	Hypertrophy of tonsils with adenoids
ICD-10-CM Codes	<p><b>J35.3</b> Hypertrophy of tonsils with hypertrophy of adenoids                      * Use additional code to identify:                      Exposure to environmental tobacco smoke (<b>Z77.22</b>)                      Exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)                      History of tobacco use (<b>Z87.891</b>)                      Occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)                      Tobacco dependence (<b>F17-</b>)                      Tobacco use (<b>Z72.0</b>)</p>

ICD-9-CM Code	ICD-10-CM Code(s)
<b>786.05</b>	Shortness of breath <b>R06.02</b> Shortness of breath
<b>493.90</b>	Asthma, unspecified
ICD-10-CM Codes	<p><b>J45.909</b> Unspecified asthma, uncomplicated                      **Use additional code to identify:                      exposure to environmental tobacco smoke (<b>Z77.22</b>)                      exposure to tobacco smoke in the perinatal period (<b>P96.81</b>)                      history of tobacco use (<b>Z87.891</b>)                      occupational exposure to environmental tobacco smoke (<b>Z57.31</b>)                      tobacco dependence (<b>F17-</b>)                      tobacco use (<b>Z72.0</b>)                      ** There are more specific code choice selections available in ICD-10-CM. These include:  <b>J45.20</b> Mild intermittent asthma, uncomplicated  <b>J45.21</b> Mild intermittent asthma with (acute) exacerbation  <b>J45.22</b> Mild intermittent asthma with status asthmaticus  <b>J45.30</b> Mild persistent asthma, uncomplicated  <b>J45.31</b> Mild persistent asthma with (acute) exacerbation  <b>J45.32</b> Mild persistent asthma with status asthmaticus  <b>J45.40</b> Moderate persistent asthma, uncomplicated  <b>J45.41</b> Moderate persistent asthma with (acute) exacerbation  <b>J45.42</b> Moderate persistent asthma with status asthmaticus  <b>J45.50</b> Severe persistent asthma, uncomplicated  <b>J45.51</b> Severe persistent asthma with (acute) exacerbation  <b>J45.52</b> Severe persistent asthma with status asthmaticus  <b>J45.901</b> Unspecified asthma with (acute) exacerbation  <b>J45.902</b> Unspecified asthma with status asthmaticus  <b>J45.990</b> Exercise induced bronchospasm  <b>J45.991</b> Cough variant asthma  <b>J45.998</b> Other asthma</p>
<b>V72.31</b>	Routine gynecological exam
ICD-10-CM Codes	<p><b>Z01.411</b> Encounter for gynecological examination (general) (routine) with abnormal findings                      * Use additional code to identify abnormal findings  <b>Z01.419</b> Encounter for gynecological examination (general) (routine) without abnormal findings                      Instructional note under codes should be as follows:                      **Use additional code:                      for screening for human papillomavirus, if applicable (<b>Z11.51</b>)                      for screening vaginal pap smear, if applicable (<b>Z12.72</b>)                      to identify acquired absence of uterus, if applicable (<b>Z90.71-</b>)</p>
<b>786.50</b>	Chest pain, unspecified
ICD-10-CM Codes	<p><b>R07.9</b> Chest pain, unspecified                      **There are more specific code choice selections available in ICD-10-CM. These include:  <b>R07.89</b> Other chest pain  <b>R07.2</b> Precordial pain  <b>R07.1</b> Chest pain on breathing  <b>R07.81</b> Pleurodynia</p>





**ICD-9-CM Code** | **ICD-10-CM Code(s)**

<b>787.20</b>	Dysphagia, unspecified
ICD-10-CM Codes	<p><b>R13.10</b> Dysphagia, unspecified  <b>**</b> There are more specific code choice selections available in ICD-10-CM. These include:  <b>R13.11</b> Dysphagia, oral phase  <b>R13.12</b> Dysphagia, oropharyngeal phase  <b>R13.13</b> Dysphagia, pharyngeal phase  <b>R13.14</b> Dysphagia, pharyngoesophageal phase  <b>R13.19</b> Other dysphagia</p>
<b>780.79</b>	Other malaise and fatigue
ICD-10-CM Codes	<p><b>R53.1</b> Weakness  <b>R53.81</b> Other malaise  <b>R53.83</b> Other fatigue  <b>G93.3</b> Postviral fatigue syndrome  <b>R53.0</b> Neoplastic (malignant) related fatigue  <i>* Code first associated neoplasm</i></p>
<b>782.3</b>	Edema
ICD-10-CM Codes	<p><b>R60.0</b> Localized edema  <b>R60.1</b> Generalized edema  <b>R60.9</b> Edema, unspecified</p>
<b>300.00</b>	Anxiety state, unspecified
ICD-10-CM Codes	<p><b>F41.1</b> Generalized anxiety disorder  <b>F41.8</b> Other specified anxiety disorders  <b>F41.9</b> Anxiety disorder, unspecified</p>
<b>530.81</b>	Esophageal reflux
ICD-10-CM Codes	<p><b>K21.9</b> Gastro-esophageal reflux disease without esophagitis  <b>K21.0</b> Gastro-esophageal reflux disease with esophagitis</p>

**ICD-9-CM Code** | **ICD-10-CM Code(s)**

<b>285.9</b>	Anemia, unspecified	
ICD-10-CM Codes	<p><b>D64.9</b> Anemia, unspecified  <b>**</b> There are more specific code choice selections available for use. These include:  <b>D64.0</b> Hereditary sideroblastic anemia  <b>D64.1</b> Secondary sideroblastic anemia due to disease  <b>**</b> Code first underlying disease  <b>D64.2</b> Secondary sideroblastic anemia due to drugs and toxins  <b>**</b> Code first (T36-T65) to identify drug or toxin  <b>**</b> Code first poisoning due to drug or toxin, if applicable (T36-T65 with fifth or sixth character 1-4 or 6)  <i>* Use additional code for adverse effect, if applicable, to identify drug (T36-T50 with fifth or sixth character 5)</i>  <b>D64.3</b> Other sideroblastic anemia  <b>D64.81</b> Anemia due to antineoplastic chemotherapy  <b>D64.89</b> Other specified anemias</p>	
<b>780.97</b>	Altered mental status	
ICD-10-CM Codes	<p><b>R41.82</b> Altered mental status, unspecified  <b>**</b> There are more specific code choice selections available in ICD-10-CM. These include:  <b>R41.0</b> Disorientation, unspecified  <b>R41.81</b> Age-related cognitive decline  <b>R41.840</b> Attention and concentration deficit  <b>R41.89</b> Other symptoms and signs involving cognitive functions and awareness</p>	
<b>276.51</b>	Dehydration	<b>E86.0</b> Dehydration
<b>786.52</b>	Painful respiration	
ICD-10-CM Codes	<p><b>R07.1</b> Chest pain on breathing  <b>R07.81</b> Pleurodynia  <b>R07.89</b> Other chest pain</p>	