



ICD-9-CM Code	ICD-10-CM Code(s)
<b>V72.31</b>	Routine gynecological exam
ICD-10-CM Codes	<p><b>Z01.411</b> Encounter for gynecological examination (general) (routine) with abnormal findings</p> <p>* Use additional code to identify abnormal findings</p> <p><b>Z01.419</b> Encounter for gynecological examination (general) (routine) without abnormal findings</p> <p>Instructional note under codes should be as follows: **Use additional code: for screening for human papillomavirus, if applicable (<b>Z11.51</b>) for screening vaginal pap smear, if applicable (<b>Z12.72</b>) to identify acquired absence of uterus, if applicable (<b>Z90.71</b>-)</p>
<b>701.0</b>	Circumscribed scleroderma
ICD-10-CM Codes	<p><b>L94.0</b> Localized scleroderma [morphea]</p> <p><b>L90.0</b> Lichen sclerosus et atrophicus</p> <p><b>L94.1</b> Linear scleroderma</p> <p><b>L94.3</b> Sclerodactyly</p>
<b>624.9</b>	Noninflammatory disorders of the vulva and perineum, unspecified
ICD-10-CM Codes	<p><b>N90.9</b> Noninflammatory disorder of vulva and perineum, unspecified</p> <p>*There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>N90.0</b> Mild vulvar dysplasia</p> <p><b>N90.1</b> Moderate vulvar dysplasia</p> <p><b>N90.3</b> Dysplasia of vulva, unspecified</p> <p><b>N90.4</b> Leukoplakia of vulva</p> <p><b>N90.5</b> Atrophy of vulva</p> <p><b>N90.6</b> Hypertrophy of vulva</p> <p><b>N90.7</b> Vulvar cyst</p> <p><b>N90.810</b> Female genital mutilation status, unspecified</p> <p><b>N90.811</b> Female genital mutilation status, Type I status</p> <p><b>N90.812</b> Female genital mutilation status, Type II status</p> <p><b>N90.813</b> Female genital mutilation status, Type III status</p> <p><b>N90.818</b> Other female genital mutilation status</p> <p><b>N90.89</b> Other specified noninflammatory disorders of vulva and perineum</p>
<b>627.1</b>	Postmenopausal bleeding
	<b>N95.0</b> Postmenopausal bleeding
<b>625.9</b>	Female genital symptoms NOS
ICD-10-CM Codes	<p><b>N94.89</b> Other specified conditions associated with female genital organs and menstrual cycle</p> <p><b>N94.9</b> Unspecified condition associated with female genital organs and menstrual cycle</p> <p><b>R10.2</b> Pelvic and perineal pain</p> <p><b>R19.8</b> Other specified symptoms and signs involving the digestive system and abdomen</p> <p><b>R39.9</b> Unspecified symptoms and signs involving the genitourinary system</p>
<b>644.03</b>	Threatened preterm labor, antepartum (without delivery)
ICD-10-CM Codes	<p><b>O47.00</b> False labor before 37 weeks of completed gestation, unspecified trimester</p> <p><b>O47.02</b> False labor before 37 completed weeks of gestation, second trimester</p> <p><b>O47.03</b> False labor before 37 completed weeks of gestation, third trimester</p>

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes (cont.)	<p><b>O47.1</b> False labor at or after 37 completed weeks of gestation</p> <p><b>O47.9</b> False labor, unspecified</p> <p>** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>
<b>618.01</b>	Cystocele, midline
	<b>N81.11</b> Cystocele, midline
<b>626.2</b>	Excessive menstruation
ICD-10-CM Codes	<p><b>N92.0</b> Excessive and frequent menstruation with regular cycle</p> <p><b>N92.1</b> Excessive and frequent menstruation with irregular cycle</p> <p><b>N92.2</b> Excessive menstruation at puberty</p> <p><b>N92.4</b> Excessive bleeding in the premenopausal period</p>
<b>648.33</b>	Drug dependence, antepartum
ICD-10-CM Codes	<p><b>O99.320</b> Drug use complicating pregnancy, unspecified trimester</p> <p><b>O99.321</b> Drug use complicating pregnancy, first trimester</p> <p><b>O99.322</b> Drug use complicating pregnancy, second trimester</p> <p><b>O99.323</b> Drug use complicating pregnancy, third trimester</p> <p>** The codes from this subcategory require an additional code from <b>F11-F16</b> and <b>F18-F19</b> to identify manifestations of the drug use.</p> <p>** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>
<b>627.3</b>	Postmenopausal atrophic vaginitis
	<b>N95.2</b> Postmenopausal atrophic vaginitis
<b>646.83</b>	Other specified complications of pregnancy, antepartum
ICD-10-CM Codes	<p><b>O26.891</b> Other specified pregnancy related conditions, first trimester</p> <p><b>O26.892</b> Other specified pregnancy related conditions, second trimester</p> <p><b>O26.893</b> Other specified pregnancy related conditions, third trimester</p> <p><b>O26.899</b> Other specified pregnancy related conditions, unspecified trimester</p> <p>* There are other code selections available for codes previously coded to <b>646.83</b>:</p> <p><b>O26.10</b> Low weight gain in pregnancy, unspecified trimester</p> <p><b>O26.11</b> Low weight gain in pregnancy, first trimester</p> <p><b>O26.12</b> Low weight gain in pregnancy, second trimester</p> <p><b>O26.13</b> Low weight gain in pregnancy, third trimester</p> <p><b>O26.40</b> Herpes gestationis, unspecified trimester</p> <p><b>O26.41</b> Herpes gestationis, first trimester</p> <p><b>O26.42</b> Herpes gestationis, second trimester</p> <p><b>O26.43</b> Herpes gestationis, third trimester</p> <p><b>O26.811</b> Pregnancy related exhaustion and fatigue, first trimester</p> <p><b>O26.812</b> Pregnancy related exhaustion and fatigue, second trimester</p> <p><b>O26.813</b> Pregnancy related exhaustion and fatigue, third trimester</p> <p><b>O26.819</b> Pregnancy related exhaustion and fatigue, unspecified trimester</p> <p>**An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>
<b>632</b>	Missed abortion
ICD-10-CM Codes	<p><b>O02.1</b> Missed abortion</p> <p>** Use additional code from category <b>O08</b> to identify any associated complication</p> <p>** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>



ICD-9-CM Code	ICD-10-CM Code(s)
<b>616.10</b>	Vaginitis and vulvovaginitis, unspecified
ICD-10-CM Codes	<p><b>N76.0</b> Acute vaginitis                      **Use additional code (<b>B95-B97</b>), to identify infectious agent                      There are other codes from this subcategory to describe vaginitis:  <b>N76.1</b> Subacute and chronic vaginitis  <b>N76.2</b> Acute vulvitis  <b>N76.3</b> Subacute and chronic vulvitis  <b>N76.89</b> Other specified inflammation of vagina and vulva</p>
<b>618.00</b>	Unspecified prolapse of vaginal walls
ICD-10-CM Codes	<p><b>N81.9</b> Female genital prolapse,unspecified                      *There are more specific code choice selections available in ICD-10-CM. These include:  <b>N81.0</b> Urethrocele  <b>N81.10</b> Cystocele, unspecified  <b>N81.11</b> Cystocele, midline  <b>N81.12</b> Cystocele, lateral  <b>N81.2</b> Incomplete uterovaginal prolapse  <b>N81.3</b> Complete uterovaginal prolapsed  <b>N81.4</b> Uterovaginal prolapsed, unspecified  <b>N81.5</b> Vaginal enterocele  <b>N81.6</b> Rectocele  <b>N81.81</b> Perineocele  <b>N81.82</b> Incompetence or weakening of pubocervical tissue  <b>N81.83</b> Incompetence or weakening of rectovaginal tissue  <b>N81.84</b> Pelvic muscle wasting  <b>N81.85</b> Cervical stump prolapsed  <b>N81.89</b> Other female genital prolapsed</p>
<b>620.2</b>	Ovarian cyst NOS/NEC
ICD-10-CM Codes	<p><b>N83.20</b> Unspecified ovarian cysts  <b>N83.29</b> Other ovarian cysts                      * There are more specific code choice selections available in ICD-10-CM. These include:  <b>N83.1</b> Corpus luteum cyst  <b>N83.0</b> Follicular cyst of ovary  <b>D27.9</b> Benign neoplasm of unspecified ovary  <b>Q50.1</b> Developmental ovarian cyst  <b>O34.80</b> Maternal care for other abnormalities of pelvic organs, unspecified trimester  <b>O34.81</b> Maternal care for other abnormalities of pelvic organs, first trimester  <b>O34.82</b> Maternal care for other abnormalities of pelvic organs, second trimester  <b>O34.83</b> Maternal care for other abnormalities of pelvic organs, third trimester  <b>O65.5</b> Obstructed labor due to abnormality of maternal pelvic organs                      ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>
<b>623.9</b>	Noninflammatory disorders of vagina, NOS
ICD-10-CM Codes	<p><b>N89.8</b> Other specified noninflammatory disorders of vagina                      * There are more specific code choice selections available in ICD-10-CM. These include:  <b>N89.0</b> Mild vaginal dysplasia  <b>N89.1</b> Moderate vaginal dysplasia  <b>N89.3</b> Dysplasia of vagina, unspecified</p>

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes (cont.)	<p><b>N89.4</b> Leukoplakia of vagina  <b>N89.5</b> Stricture and atresia of vagina  <b>N89.6</b> Tight hymenal ring  <b>N89.7</b> Hematocolpos  <b>N89.9</b> Noninflammatory disorder of vagina, unspecified</p>
<b>V25.2</b>	Desire for sterilization <b>Z30.2</b> Encounter for sterilization
<b>618.1</b>	Uterine Prolapse w/o mention of vaginal wall prolapse
ICD-10-CM Codes	<p><b>N81.2</b> Incomplete uterovaginal prolapse  <b>N81.3</b> Complete uterovaginal prolapse  <b>N81.4</b> Uterovaginal prolapse, unspecified</p>
<b>788.20</b>	Urinary retention, NOS
ICD-10-CM Codes	<p><b>R33.9</b> Retention of urine, unspecified                      * There are more specific code choice selections available in ICD-10-CM. These include:  <b>R33.8</b> Other retention of urine  <b>R33.0</b> Drug induced retention of urine                      ** Use additional code for adverse effect, if applicable, to identify drug (<b>T36-T50</b> with fifth or sixth character 5)  <b>F45.8</b> Other somatoform disorders</p>
<b>788.31</b>	Urge incontinence
ICD-10-CM Codes	<p><b>N39.41</b> Urge incontinence                      ** Code also any associated overactive bladder (<b>N32.81</b>)</p>
<b>788.33</b>	Mixed incontinence
ICD-10-CM Codes	<p><b>N39.46</b> Mixed incontinence                      ** Code also any associated overactive bladder symptoms (<b>N32.81</b>)</p>
<b>789.03</b>	Abdominal pain RLQ <b>R10.31</b> Right lower quadrant pain
<b>625.6</b>	Female stress incontinence
ICD-10-CM Codes	<p><b>N39.3</b> Stress incontinence (female) (male)                      ** Code also any associated overactive bladder symptoms (<b>N32.81</b>)</p>
<b>V22.1</b>	Supervision other normal pregnancy
ICD-10-CM Codes	<p><b>Z34.80</b> Encounter for supervision of other normal pregnancy, unspecified trimester  <b>Z34.81</b> Encounter for supervision of other normal pregnancy, first trimester  <b>Z34.82</b> Encounter for supervision of other normal pregnancy, second trimester  <b>Z34.83</b> Encounter for supervision of other normal pregnancy, third trimester</p>
<b>611.72</b>	Lump or mass in breast <b>N63</b> Unspecified lump in breast
<b>599.0</b>	Urinary tract infection, NOS
ICD-10-CM Codes	<p><b>N39.0</b> Urinary tract infection, site not specified                      ** Use additional code (<b>B95-B97</b>), to identify infectious agent                      *There are more specific code choice selections available in ICD-10-CM. These include:  <b>N30.00</b> Acute cystitis without hematuria  <b>N30.01</b> Acute cystitis with hematuria</p>



ICD-9-CM Code	ICD-10-CM Code(s)
<b>599.0</b>	Continued
ICD-10-CM Codes (cont.)	<p><b>N30.10</b> Interstitial cystitis (chronic) without hematuria  <b>N30.11</b> Interstitial cystitis (chronic) with hematuria  <b>N30.20</b> Other chronic cystitis without hematuria  <b>N30.21</b> Other chronic cystitis with hematuria  <b>N30.30</b> Trigonitis without hematuria  <b>N30.31</b> Trigonitis with hematuria  <b>N30.40</b> Irradiation cystitis without hematuria  <b>N30.41</b> Irradiation cystitis with hematuria  <b>N30.80</b> Other cystitis without hematuria  <b>N30.81</b> Other cystitis with hematuria  <b>N30.90</b> Cystitis, unspecified without hematuria  <b>N30.91</b> Cystitis, unspecified with hematuria  <b>N15.9</b> Renal tubule-interstitial disease, unspecified  <b>N34.1</b> Nonspecific urethritis  <b>N34.2</b> Other urethritis</p>
<b>627.9</b>	Menopausal disorder, NOS
ICD-10-CM Codes	<p><b>N95.9</b> Unspecified menopausal and perimenopausal disorder  *There are more specific code choice selections available in ICD-10-CM. These include:  <b>N95.0</b> Postmenopausal bleeding  <b>N95.1</b> Menopausal and female climacteric states  <b>N95.2</b> Postmenopausal atrophic vaginitis  <b>N95.8</b> Other specified menopausal and perimenopausal disorders</p>
<b>788.34</b>	Incontinence without sensory awareness
ICD-10-CM Codes	<p><b>N39.42</b> Incontinence without sensory awareness  ** Code also any associated overactive bladder (<b>N32.81</b>)</p>
<b>642.43</b>	Mild or unspecified pre-eclampsia, NOS
ICD-10-CM Codes	<p><b>O14.00</b> Mild to moderate pre-eclampsia, unspecified trimester  <b>O14.02</b> Mild to moderate pre-eclampsia, second trimester  <b>O14.03</b> Mild to moderate pre-eclampsia, third trimester  <b>O14.90</b> Unspecified pre-eclampsia, unspecified trimester  <b>O14.92</b> Unspecified pre-eclampsia, second trimester  <b>O14.93</b> Unspecified pre-eclampsia, third trimester  **An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>
<b>795.01</b>	Papanicolaou smear of cervix with atypical squamous cells of undetermined significance ASC-US
ICD-10-CM Codes	<p><b>R87.610</b> Atypical squamous cells of undetermined significance on cytologic smear of cervix (ASC-US)  *There are more specific code choice selections available in ICD-10-CM. These include:  <b>R87.611</b> Atypical squamous cells cannot exclude high grade squamous intraepithelial lesion on cytologic smear of cervix (ASC-H)  <b>R87.612</b> Low grade squamous intraepithelial lesion on cytologic smear of cervix (LGSIL)  <b>R87.613</b> High grade squamous intraepithelial lesion on cytologic smear of cervix (HGSIL)  <b>R87.614</b> Cytologic evidence of malignancy on smear of cervix  <b>R87.615</b> Unsatisfactory cytologic smear of cervix</p>

ICD-9-CM Code	ICD-10-CM Code(s)
<b>795.01</b>	Continued
ICD-10-CM Codes (cont.)	<p><b>R87.616</b> Satisfactory cervical smear but lacking transformation zone  <b>R87.618</b> Other abnormal cytological findings on specimens from cervix uteri  <b>R87.619</b> Unspecified abnormal cytological findings in specimens from cervix uteri</p>
<b>626.4</b>	Irregular menstrual cycle
ICD-10-CM Codes	<p><b>N92.5</b> Other specified irregular menstruation  <b>N92.6</b> Irregular menstruation, unspecified  * There are more specific code choice selections available in ICD-10-CM. These include:  <b>N92.0</b> Excessive and frequent menstruation with regular cycle  <b>N92.1</b> Excessive and frequent menstruation with irregular cycle  <b>N92.2</b> Excessive menstruation at puberty  <b>N92.4</b> Excessive bleeding in the premenopausal period</p>
<b>651.03</b>	Twin pregnancy antepartum
ICD-10-CM Codes	<p><b>O30.001</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, first trimester  <b>O30.002</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, second trimester  <b>O30.003</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, third trimester  <b>O30.009</b> Twin pregnancy, unspecified number of placenta and unspecified number of amniotic sacs, unspecified trimester  <b>O30.011</b> Twin pregnancy, monochorionic/monoamniotic, first trimester  <b>O30.012</b> Twin pregnancy, monochorionic/monoamniotic, second trimester  <b>O30.013</b> Twin pregnancy, monochorionic/monoamniotic, third trimester  <b>O30.019</b> Twin pregnancy, monochorionic/monoamniotic, unspecified trimester  <b>O30.031</b> Twin pregnancy, monochorionic/diamniotic, first trimester  <b>O30.032</b> Twin pregnancy, monochorionic/diamniotic, second trimester  <b>O30.033</b> Twin pregnancy, monochorionic/diamniotic, third trimester  <b>O30.039</b> Twin pregnancy, monochorionic/diamniotic, unspecified trimester  <b>O30.041</b> Twin pregnancy, dichorionic/diamniotic, first trimester  <b>O30.042</b> Twin pregnancy, dichorionic/diamniotic, second trimester  <b>O30.043</b> Twin pregnancy, dichorionic/diamniotic, third trimester  <b>O30.049</b> Twin pregnancy, dichorionic/diamniotic, unspecified trimester  <b>O30.091</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, first trimester  <b>O30.092</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, second trimester  <b>O30.093</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, third trimester  <b>O30.099</b> Twin pregnancy, unable to determine number of placenta and number of amniotic sacs, unspecified trimester  **An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>



ICD-9-CM Code	ICD-10-CM Code(s)	
<b>795.03</b>	Papanicolaou smear of cervix with LGSIL	
ICD-10-CM Codes	<b>R87.612</b> Low grade squamous intraepithelial lesion on cytologic smear of cervix (LGSIL)	
<b>623.5</b>	Leukorrhea, not specified as infective	<b>N89.8</b> Other specified noninflammatory disorders of vagina
<b>V25.11</b>	Insertion of IUD	<b>Z30.430</b> Encounter for insertion of intrauterine contraceptive device
<b>658.23</b>	Delayed delivery after spontaneous or unspecified rupture of membranes, antepartum condition or complication	
ICD-10-CM Codes	<p><b>O42.111</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, first trimester</p> <p><b>O42.112</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, second trimester</p> <p><b>O42.113</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, third trimester</p> <p><b>O42.119</b> Preterm premature rupture of membranes, onset of labor more than 24 hours following rupture, unspecified trimester</p> <p>** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>	
<b>789.00</b>	Abdominal pain site unspecified	
ICD-10-CM Codes	<p><b>R10.9</b> Unspecified abdominal pain * There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>R10.0</b> Acute abdomen</p> <p><b>R10.10</b> Upper abdominal pain, unspecified</p> <p><b>R10.11</b> Right upper quadrant pain</p> <p><b>R10.12</b> Left upper quadrant pain</p> <p><b>R10.13</b> Epigastric pain</p> <p><b>R10.2</b> Pelvic and perineal pain</p> <p><b>R10.30</b> Lower abdominal pain, unspecified</p> <p><b>R10.31</b> Right lower quadrant pain</p> <p><b>R10.32</b> Left lower quadrant pain</p> <p><b>R10.33</b> Periumbilical pain</p> <p><b>R10.84</b> Generalized abdominal pain</p>	
<b>V28.3</b>	Encounter for routine screening for malformation using ultrasonics (routine fetal ultrasound, NOS)	<b>Z36</b> Encounter for antenatal screening of mother
<b>V76.2</b>	Screening malignant neoplasm cervix	<b>Z12.4</b> Encounter for screening for malignant neoplasm of cervix
<b>655.13</b>	Maternal care for fetal chromosomal abnormality	
ICD-10-CM Codes	<p>*A 7th character extender must be assigned to code under category <b>O35</b>.</p> <p><b>O35.1XX0</b> Maternal care for (suspected) chromosomal abnormality in fetus, not applicable or unspecified</p> <p><b>O35.1XX1</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 1</p> <p><b>O35.1XX2</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 2</p>	

ICD-9-CM Code	ICD-10-CM Code(s)	
ICD-10-CM Codes (cont.)	<p><b>O35.1XX3</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 3</p> <p><b>O35.1XX4</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 4</p> <p><b>O35.1XX5</b> Maternal care for (suspected) chromosomal abnormality in fetus, fetus 5</p> <p><b>O35.1XX9</b> Maternal care for (suspected) chromosomal abnormality in fetus, other fetus</p> <p>One of the following 7th characters is to be assigned to each code under category O35. 7th character 0 is for single gestations and multiple gestations where the fetus is unspecified. 7th character 1 through 9 are for cases of multiple gestations to identify the fetus for which the code applies. The appropriate code from category O30, multiple gestation, must also be assigned when assigning a code from category O35 that has a 7th character of 1 through 9.</p> <p>7th Character extenders for subcategory <b>O35-</b></p> <p><b>0</b> Not Applicable or unspecified</p> <p><b>1</b> Fetus 1</p> <p><b>2</b> Fetus 2</p> <p><b>3</b> Fetus 3</p> <p><b>4</b> Fetus 4</p> <p><b>5</b> Fetus 5</p> <p><b>9</b> Other fetus</p> <p>**An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.</p>	
<b>789.04</b>	Abdominal pain LLQ	<b>R10.32</b> Left lower quadrant pain
<b>789.07</b>	Abdominal pain generalized	<b>R10.84</b> Generalized abdominal pain
<b>595.1</b>	Interstitial cystitis chronic	
ICD-10-CM Codes	<p><b>N30.10</b> Interstitial cystitis (chronic) without hematuria</p> <p><b>N30.11</b> Interstitial cystitis (chronic) with hematuria</p>	
<b>595.9</b>	Cystitis NOS	
ICD-10-CM Codes	<p><b>N30.90</b> Cystitis, unspecified without hematuria</p> <p><b>N30.91</b> Cystitis, unspecified with hematuria</p> <p>**Use additional code to identify infectious agent (<b>B95-B97</b>)</p> <p>* There are more specific code choice selections available in ICD-10-CM. These include:</p> <p><b>N30.00</b> Acute cystitis without hematuria</p> <p><b>N30.01</b> Acute cystitis with hematuria</p> <p><b>N30.10</b> Interstitial cystitis (chronic) without hematuria</p> <p><b>N30.11</b> Interstitial cystitis (chronic) with hematuria</p> <p><b>N30.20</b> Other chronic cystitis without hematuria</p> <p><b>N30.21</b> Other chronic cystitis with hematuria</p> <p><b>N30.30</b> Trigonitis without hematuria</p> <p><b>N30.31</b> Trigonitis with hematuria</p> <p><b>N30.40</b> Irradiation cystitis without hematuria</p> <p><b>N30.41</b> Irradiation cystitis with hematuria</p> <p><b>N30.80</b> Other cystitis without hematuria</p> <p><b>N30.81</b> Other cystitis with hematuria</p>	
<b>788.43</b>	Nocturia	<b>R35.1</b> Nocturia
<b>112.1</b>	Candidial vaginitis	<b>B37.3</b> Candidiasis of vulva and vagina
<b>616.2</b>	Bartholin's gland cyst	<b>N75.0</b> Cyst of Bartholin's gland



ICD-9-CM Code	ICD-10-CM Code(s)
<b>616.4</b>	Abcess of vulva NEC <b>N76.4</b> Abscess of vulva
<b>V25.09</b>	Contraceptive management NEC
ICD-10-CM Codes	<b>Z30.09</b> Encounter for other general counseling and advice on contraception <b>Z30.8</b> Encounter for other contraceptive management
<b>659.73</b>	Abnormality in fetal heart rate or rhythm, antepartum
ICD-10-CM Codes	<b>O76</b> Abnormality in fetal heart rate and rhythm complicating labor and delivery ** An additional code from category <b>Z3A</b> is needed to specify the weeks of gestation of the pregnancy.
<p>Many codes from chapter 15 (<b>O00-O99A</b>) require the use of an additional code from category <b>Z3A</b>. Weeks of gestation, to identify the specific week of the pregnancy. These codes are for use, only on the maternal record, to indicate the weeks of gestation of the pregnancy.</p>	
ICD-10-CM Codes	Code first complications of pregnancy, childbirth and the puerperium ( <b>O00-O9A</b> ). <b>Z3A.00</b> Weeks of gestation of pregnancy not specified <b>Z3A.01</b> Less than 8 weeks gestation of pregnancy <b>Z3A.08</b> 8 weeks gestation of pregnancy <b>Z3A.09</b> 9 weeks gestation of pregnancy <b>Z3A.10</b> 10 weeks gestation of pregnancy <b>Z3A.11</b> 11 weeks gestation of pregnancy <b>Z3A.12</b> 12 weeks gestation of pregnancy

ICD-9-CM Code	ICD-10-CM Code(s)
ICD-10-CM Codes	<b>Z3A.13</b> 13 weeks gestation of pregnancy <b>Z3A.14</b> 14 weeks gestation of pregnancy <b>Z3A.15</b> 15 weeks gestation of pregnancy <b>Z3A.16</b> 16 weeks gestation of pregnancy <b>Z3A.17</b> 17 weeks gestation of pregnancy <b>Z3A.18</b> 18 weeks gestation of pregnancy <b>Z3A.19</b> 19 weeks gestation of pregnancy <b>Z3A.20</b> 20 weeks gestation of pregnancy <b>Z3A.21</b> 21 weeks gestation of pregnancy <b>Z3A.22</b> 22 weeks gestation of pregnancy <b>Z3A.23</b> 23 weeks gestation of pregnancy <b>Z3A.24</b> 24 weeks gestation of pregnancy <b>Z3A.25</b> 25 weeks gestation of pregnancy <b>Z3A.26</b> 26 weeks gestation of pregnancy <b>Z3A.27</b> 27 weeks gestation of pregnancy <b>Z3A.28</b> 28 weeks gestation of pregnancy <b>Z3A.29</b> 29 weeks gestation of pregnancy <b>Z3A.30</b> 30 weeks gestation of pregnancy <b>Z3A.31</b> 31 weeks gestation of pregnancy <b>Z3A.32</b> 32 weeks gestation of pregnancy <b>Z3A.33</b> 33 weeks gestation of pregnancy <b>Z3A.34</b> 34 weeks gestation of pregnancy <b>Z3A.35</b> 35 weeks gestation of pregnancy <b>Z3A.36</b> 36 weeks gestation of pregnancy <b>Z3A.37</b> 37 weeks gestation of pregnancy <b>Z3A.38</b> 38 weeks gestation of pregnancy <b>Z3A.39</b> 39 weeks gestation of pregnancy <b>Z3A.40</b> 40 weeks gestation of pregnancy <b>Z3A.41</b> 41 weeks gestation of pregnancy <b>Z3A.42</b> 42 weeks gestation of pregnancy <b>Z3A.49</b> Greater than 42 weeks gestation of pregnancy

# Fast Forward

to ICD-10



# *Fast Forward*

*to ICD-10*

