

ICD-9-CM Code	ICD-10-CM Code(s)
185	Malignant neoplasm of prostate C61 Malignant neoplasm of prostate
600.01	Hypertrophy (benign) of prostate with urinary obstruction and other lower urinary tract symptoms (LUTS) lower urinary tract symptoms (LUTS)
ICD-10-CM Codes	Two codes are required in ICD-10-CM: N40.1 Enlarged prostate with lower urinary tract symptoms (LUTS); AND N13.8 Other obstructive and reflux uropathy ** Note under N40.1 states – Use additional code for associated symptoms, when specified: <ul style="list-style-type: none"> • Incomplete bladder emptying (R39.14) • Nocturia (R35.1) • Straining on urination (R39.16) • Urinary frequency (R35.0) • Urinary hesitancy (R39.11) • Urinary incontinence (N39.4-) • Urinary obstruction (N13.8) • Urinary retention (R33.8) • Urinary urgency (R39.15) • Weak urinary stream (R39.12) ** Note under N13.8 states – Code first, if applicable, any causal condition, such as: <ul style="list-style-type: none"> • Enlarged prostate (N40.1)
788.20	Retention of urine, unspecified
ICD-10-CM Codes	R33.9 Retention of urine, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: R33.0 Drug induced retention of urine ** Use additional code for adverse effect, if applicable, to identify drug (T36-T50 with fifth or sixth character 5) R33.8 Other retention of urine ** Code first, if applicable, any causal condition, such as: enlarged prostate (N40.1)
188.9	Malignant neoplasm of bladder, part unspecified
ICD-10-CM Codes	C67.9 Malignant neoplasm of bladder, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: C67.0 Malignant neoplasm of trigone of bladder C67.1 Malignant neoplasm of dome of bladder C67.2 Malignant neoplasm of lateral wall of bladder C67.3 Malignant neoplasm of anterior wall of bladder C67.4 Malignant neoplasm of posterior wall of bladder C67.5 Malignant neoplasm of bladder neck C67.6 Malignant neoplasm of ureteric orifice C67.7 Malignant neoplasm of urachus C67.8 Malignant neoplasm of overlapping sites of bladder
607.84	Impotence of organic origin
ICD-10-CM Codes	N52.9 Male erectile dysfunction, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: N52.01 Erectile dysfunction due to arterial insufficiency N52.02 Corporo-venous occlusive erectile dysfunction N52.03 Combined arterial insufficiency and corporo-venous occlusive erectile dysfunction

ICD-9-CM Code	ICD-10-CM Code(s)
607.84	Continued
ICD-10-CM Codes (cont.)	N52.1 Erectile dysfunction due to diseases classified elsewhere ** Code first underlying disease N52.2 Drug-induced erectile dysfunction N52.31 Erectile dysfunction following radical prostatectomy N52.32 Erectile dysfunction following radical cystectomy N52.33 Erectile dysfunction following urethral surgery N52.34 Erectile dysfunction following simple prostatectomy N52.39 Other post-surgical erectile dysfunction N52.8 Other male erectile dysfunction
592.0	Calculus of kidney
ICD-10-CM Codes	N20.0 Calculus of kidney N20.9 Urinary calculus; unspecified N20.2 Calculus of kidney and ureter
790.93	Elevated prostate specific antigen (PSA) R97.2 Elevated prostate specific antigen (PSA)
788.21	Incomplete bladder emptying N20.2 Calculus of kidney with calculus of ureter
788.31	Urge incontinence N39.41 Urge incontinence
592.1	Calculus of ureter
ICD-10-CM Codes	N20.1 Calculus of ureter N20.2 Calculus of kidney with calculus of ureter
596.54	Neurogenic bladder NOS
ICD-10-CM Codes	N31.9 Neuromuscular dysfunction of bladder, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: N31.0 Uninhibited neuropathic bladder, not elsewhere classified N31.1 Reflex neuropathic bladder, not elsewhere classified N31.2 Flaccid neuropathic bladder, not elsewhere classified N31.8 Other neuromuscular dysfunction of bladder ** Use additional code to identify any associated urinary incontinence (N39.3-N39.4-)
189.0	Malignant neoplasm of kidney, except pelvis
ICD-10-CM Codes	C64.1 Malignant neoplasm of right kidney, except renal pelvis C64.2 Malignant neoplasm of left kidney, except renal pelvis C64.9 Malignant neoplasm of unspecified kidney, except renal pelvis
788.41	Urinary frequency R35.0 Frequency of micturition ** Code first, if applicable, any causal condition, such as enlarged prostate
599.0	Urinary tract infection, site not specified
ICD-10-CM Codes	N39.0 Urinary tract infection, site not specified ** Use additional code (B95-B97) to identify infectious agent

ICD-9-CM Code	ICD-10-CM Code(s)
599.0	Continued
ICD-10-CM Codes (cont.)	<p>** There are more specific code choice selections available in ICD-10-CM. These include:</p> <p>N30.00 Acute cystitis without hematuria N30.01 Acute cystitis with hematuria N30.10 Interstitial cystitis (chronic) without hematuria N30.11 Interstitial cystitis (chronic) with hematuria N30.20 Other chronic cystitis without hematuria N30.21 Other chronic cystitis with hematuria N30.30 Trigonitis without hematuria N30.31 Trigonitis with hematuria N30.40 Irradiation cystitis without hematuria N30.41 Irradiation cystitis with hematuria N30.80 Other cystitis without hematuria N30.81 Other cystitis with hematuria N30.90 Cystitis, unspecified without hematuria N30.91 Cystitis, unspecified with hematuria N15.9 Renal tubule-interstitial disease, unspecified N34.1 Nonspecific urethritis N34.2 Other urethritis</p>
625.6	Stress incontinence, female
ICD-10-CM Codes	<p>N39.3 Stress incontinence (female) (male) ** Code also any associated overactive bladder (N32.81).</p>
599.71	Gross hematuria
ICD-10-CM Codes	<p>R31.0 Gross hematuria R31.1 Benign essential microscopic hematuria R31.2 Other microscopic hematuria R31.9 Unspecified hematuria</p>
788.32	Stress incontinence, male
ICD-10-CM Codes	<p>N39.3 Stress incontinence (female) (male) ** Code also any associated overactive bladder (N32.81).</p>
239.5	Neoplasm of unspecified nature, other genitourinary organs
	D49.5 Neoplasm of unspecified behavior of other genitourinary organs
598.9	Urethral stricture, unspecified
ICD-10-CM Codes	<p>N35.9 Urethral stricture, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: N35.010 Post-traumatic urethral stricture, male, meatal N35.011 Post-traumatic bulbous urethral stricture N35.012 Post-traumatic membranous urethral stricture N35.013 Post-traumatic anterior urethral stricture N35.014 Post-traumatic urethral stricture, male, unspecified N35.021 Urethral stricture due to childbirth N35.028 Other post-traumatic urethral stricture, female N35.111 Postinfective urethral stricture, not elsewhere classified, male, meatal N35.112 Postinfective bulbous urethral stricture, not elsewhere classified</p>

ICD-9-CM Code	ICD-10-CM Code(s)
598.9	Continued
ICD-10-CM Codes (cont.)	<p>N35.113 Postinfective membranous urethral stricture, not elsewhere classified N35.114 Postinfective anterior urethral stricture, not elsewhere classified N35.119 Postinfective urethral stricture, not elsewhere classified, male, unspecified N35.12 Postinfective urethral stricture, not elsewhere classified, female N35.8 Other urethral stricture, unspecified</p>
605	Redundant prepuce and phimosis
ICD-10-CM Codes	<p>N47.0 Adherent prepuce, newborn N47.1 Phimosis N47.2 Paraphimosis N47.8 Other disorders of prepuce</p>
591	Hydronephrosis
ICD-10-CM Codes	<p>N13.30 Unspecified hydronephrosis ** There are more specific code choice selections available in ICD-10-CM. These include: N13.1 Hydronephrosis with ureteral stricture, not elsewhere classified N13.2 Hydronephrosis with renal and ureteral calculus obstruction N13.39 Other hydronephrosis</p>
603.9	Hydrocele, unspecified
ICD-10-CM Codes	<p>N43.3 Hydrocele, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: N43.0 Encysted hydrocele N43.1 Infected hydrocele ** Use additional code (B95-B97) to identify infectious agent N43.2 Other hydrocele</p>
239.4	Neoplasm of unspecified nature, bladder
	D49.4 Neoplasm of unspecified behavior of bladder
788.43	Nocturia
	R35.1 Nocturia
257.2	Other testicular hypofunction
	E29.1 Testicular hypofunction **Use additional code for adverse effect, if applicable, to identify drug (T36-T50 with fifth or sixth character 5)
600.91	Hyperplasia of prostate, unspecified, with urinary obstruction and other lower urinary tract symptoms (LUTS)
ICD-10-CM Codes	<p>Two codes are required in ICD-10-CM: N40.1 Enlarged prostate with lower urinary tract symptoms (LUTS); AND N13.8 Other obstructive and reflux uropathy *** ** Note under N40.1 states – Use additional code for associated symptoms, when specified: <ul style="list-style-type: none"> • Incomplete bladder emptying (R39.14) • Nocturia (R35.1) • Straining on urination (R39.16) </p>



ICD-9-CM Code	ICD-10-CM Code(s)	
600.91 (cont.)	Hyperplasia of prostate, unspecified, with urinary obstruction and other lower urinary tract symptoms (LUTS)	
ICD-10-CM Codes (cont.)	<ul style="list-style-type: none"> • Urinary frequency (R35.0) • Urinary hesitancy (R39.11) • Urinary incontinence (N39.4-) • Urinary obstruction (N13.8) • Urinary retention (R33.8) • Urinary urgency (R39.15) • Weak urinary stream (R39.12) <p>** Note under N13.8 states – Code first, if applicable, any causal condition, such as:</p> <ul style="list-style-type: none"> • Enlarged prostate (N40.1) 	
594.1	Other calculus in bladder	N21.0 Calculus in bladder
595.2	Other chronic cystitis	
ICD-10-CM Codes	N30.20 Other chronic cystitis without hematuria N30.21 Other chronic cystitis with hematuria ** Use additional code to identify infectious agent (B95-B97).	
601.1	Chronic prostatitis	N41.1 Chronic prostatitis
600.00	Hypertrophy (benign) of prostate without urinary obstruction and other lower urinary tract symptoms (LUTS)	N40.0 Enlarged prostate without lower urinary tract symptoms (LUTS)
752.51	Undescended testis	
ICD-10-CM Codes	Q53.10 Unspecified undescended testicle, unilateral Q53.11 Abdominal testicle, unilateral Q53.12 Ectopic perineal testis, unilateral Q53.20 Undescended testicle, unspecified, bilateral Q53.21 Abdominal testis, bilateral Q53.22 Ectopic perineal testis, bilateral Q53.9 Hypospadias, unspecified	
788.30	Urinary incontinence, unspecified	
ICD-10-CM Codes	R32 Unspecified urinary incontinence R39.81 Functional urinary incontinence F98.0 Non-organic origin ** There are more specific code choice selections available in ICD-10-CM. These include: N39.3 Stress incontinence (female) (male) ** Code also associated overactive bladder (N32.81) N39.41 Urge incontinence N39.42 Incontinence without sensory awareness N39.43 Post-void dribbling N39.44 Nocturnal enuresis N39.45 Continuous leakage N39.46 Mixed incontinence N39.490 Overflow incontinence N39.498 Other specified urinary incontinence ** Code also any associated overactive bladder (N32.81)	

ICD-9-CM Code	ICD-10-CM Code(s)	
595.1	Chronic interstitial cystitis	
ICD-10-CM Codes	N30.10 Interstitial cystitis (chronic) without hematuria ** N30.11 Interstitial cystitis (chronic) with hematuria ** ** Use additional code to identify infectious agent (B95-B97).	
593.4	Other ureteric obstruction	
ICD-10-CM Codes	N13.8 Other obstructive and reflux uropathy ** Code first, if applicable, any causal condition, such as: enlarged prostate (N40.1).	
585.9	Chronic kidney disease, unspecified	
ICD-10-CM Codes	N18.9 Chronic kidney disease, unspecified ** Code first any associated: <ul style="list-style-type: none"> • Diabetic chronic kidney disease (E08.22, E09.22, E10.22, E11.22, E13.22) • Hypertensive chronic kidney disease (I12-, I13-) • Use additional code to identify kidney transplant status, if applicable (Z94.0) ** There are more specific code choice selections available in ICD-10-CM. These include: N18.1 Chronic kidney disease, stage 1 N18.2 Chronic kidney disease, stage 2 (mild) N18.3 Chronic kidney disease, stage 3 (moderate) N18.4 Chronic kidney disease, stage 4 (severe) N18.5 Chronic kidney disease, stage 5 N18.6 End stage renal disease ** Use additional code to identify dialysis status (Z99.2)	
788.63	Urgency of urination	
ICD-10-CM Codes	R39.15 Urgency of urination ** Code first, if applicable, any causal condition, such as enlarged prostate (N40.1).	
788.1	Dysuria	R30.0 Dysuria
599.70	Hematuria, unspecified	
ICD-10-CM Codes	R31.9 Hematuria, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: R31.0 Gross hematuria R31.1 Benign essential microscopic hematuria R31.2 Other microscopic hematuria N02.0 Recurrent and persistent hematuria with minor glomerular abnormality N02.1 Recurrent and persistent hematuria with focal and segmental glomerular lesions N02.2 Recurrent and persistent hematuria with diffuse membranous glomerulonephritis N02.3 Recurrent and persistent hematuria with diffuse mesangial proliferative glomerulonephritis N02.4 Recurrent and persistent hematuria with diffuse endocapillary proliferative glomerulonephritis N02.5 Recurrent and persistent hematuria with diffuse mesangiocapillary glomerulonephritis N02.6 Recurrent and persistent hematuria with dense deposit disease	



ICD-9-CM Code	ICD-10-CM Code(s)
599.70	Continued
ICD-10-CM Codes (cont.)	N02.7 Recurrent and persistent hematuria with diffuse crescentic glomerulonephritis N02.8 Recurrent and persistent hematuria with other morphologic changes N02.9 Recurrent and persistent hematuria with unspecified morphologic changes
593.2	Cyst of kidney, acquired N28.1 Cyst of kidney, acquired
599.72	Microscopic hematuria
ICD-10-CM Codes	R31.1 Benign essential microscopic hematuria R31.2 Other microscopic hematuria
607.1	Balanoposthitis
ICD-10-CM Codes	N47.6 Balanoposthitis N48.1 Balanitis ** Use additional code (B95-B97) to identify infectious agent.
789.00	Abdominal pain, unspecified site
ICD-10-CM Codes	R10.9 Unspecified abdominal pain ** There are more specific code choice selections available in ICD-10-CM. These include: R10.0 Acute abdomen R10.10 Upper abdominal pain, unspecified R10.11 Right upper quadrant pain R10.12 Left upper quadrant pain R10.13 Epigastric pain R10.2 Pelvic and perineal pain R10.30 Lower abdominal pain, unspecified R10.31 Right lower quadrant pain R10.32 Left lower quadrant pain R10.33 Periumbilical pain R10.84 Generalized abdominal pain
752.61	Hypospadias
ICD-10-CM Codes	Q54.9 Hypospadias, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: Q54.0 Hypospadias, balanic Q54.1 Hypospadias, penile Q54.2 Hypospadias, penoscrotal Q54.3 Hypospadias, perineal Q54.8 Other hypospadias

ICD-9-CM Code	ICD-10-CM Code(s)
596.9	Unspecified disorder of bladder
ICD-10-CM Codes	N32.9 Bladder disorder, unspecified ** There are more specific code choice selections available in ICD-10-CM. These include: N32.0 Bladder neck obstruction N32.3 Diverticulum of bladder
788.36	Nocturnal enuresis N39.44 Nocturnal enuresis
600.10	Nodular prostate without urinary obstruction N40.0 Enlarged prostate without lower urinary tract symptoms (LUTS)
608.9	Unspecified disorder of male genital organs N50.9 Disorder of male genital organs, unspecified
604.90	Orchitis and epididymitis, unspecified
ICD-10-CM Codes	N45.1 Epididymitis N45.2 Orchitis N45.3 Epididymo-orchitis ** Use additional code (B95-B97) to identify infectious agent
608.1	Spermatocele
ICD-10-CM Codes	N43.40 Spermatocele of epididymis, unspecified N43.41 Spermatocele of epididymis, single N43.42 Spermatocele of epididymis, multiple
607.85	Peyronie's disease N48.6 Peyronie's disease
598.2	Postoperative urethral stricture
ICD-10-CM Codes	N99.110 Postprocedural urethral stricture, male, meatal N99.111 Postprocedural bulbous urethral stricture N99.112 Postprocedural membranous urethral stricture N99.113 Postprocedural anterior urethral stricture N99.114 Postprocedural urethral stricture, male, unspecified N99.12 Postprocedural urethral stricture, female